



Written by [R. Cort Kirkwood](#) on March 3, 2023

CDC: Sodomy Spreading Shigellosis, a Drug-resistant Intestinal Disease

The Centers for Disease Control has issued an official health advisory that says a drug-resistant variant of the shigella bacterium is spreading. And it spreads mostly through homosexual sodomy.

The microbe causes the diarrhea-inducing shigellosis infection, although it is also contracted through contaminated food and drink.

This outbreak of the disease follows last year's monkeypox contagion, which is also spread by the same activity.

To its credit, CDC acknowledges that fact openly.



cdc.gov

CDC issues new Health Alert Network (HAN) health advisory. CDC observed an increase in extensively drug-resistant shigellosis among some populations of adults. See recommendations for diagnosing & managing these infections & steps to keep others healthy: <https://t.co/xtJpUPfoAa> pic.twitter.com/4uwpOI7oWH

— CDC (@CDCgov) [March 1, 2023](#)

The Advisory: Fecal-oral Route

But worse than the way it spreads, which is bad enough, is that a number of the infections in the outbreak are drug-resistant, or XDR.

"In 2022, about 5% of Shigella infections reported to CDC were caused by XDR strains, compared with 0% in 2015," [CDC reports](#). "Clinicians treating patients infected with XDR strains have limited antimicrobial treatment options."

Symptoms include inflammatory and sometimes bloody diarrhea, along with fever, stomach cramps, and [tenesmus](#).

Treatment doesn't typically require drugs.

"Most people with diarrheal illness require only supportive care and fluid replacement," CDC reports. Antibiotics aren't always needed, but they may be indicated to shorten the duration of illness (by about 2 days), or reduce the likelihood of transmission."

The microbe is "transmitted by the fecal-oral route, directly through person-to-person contact including sexual contact, and indirectly through contaminated food, water, and other routes," the [advisory explains](#).

Homosexual sodomy explains a change in age of those typically infected:



Written by [R. Cort Kirkwood](#) on March 3, 2023

Historically, shigellosis has predominantly affected young children (age 1–4 years) in the United States.

More recently, CDC has observed an increase in antimicrobial-resistant *Shigella* infections among adult populations.

Tops on that list are “Gay, bisexual, and other men who have sex with men.”



Indeed, the CDC’s [opening *Shigella* page asks](#) this question front and center, accompanied by a picture of two homosexuals embracing: Did you know the diarrhea germ *Shigella* can be spread through sex activity? Learn more.”

“Gay, bisexual, and other men who have sex with men* are among groups at high risk for infection with *Shigella* germs,” the [website continues](#). “*Shigella* germs spread easily and rapidly among people, including during sexual activity”:

Men who have sex with men are particularly at risk for infections with antimicrobial-resistant *Shigella*. Antimicrobial resistance happens when germs like bacteria and fungi develop the ability to defeat the drugs designed to kill them. That means the germs are not killed and continue to grow. *Shigella* germs are increasingly resistant to antibiotic treatment.

CDC also warns homosexuals to “use latex gloves during anal fingering or fisting” and to “wash sex toys with soap and water after each use and wash hands after touching used sex toys.”

And, CDC continues, “if you or your partner has diarrhea, do not have sex. To reduce the chance of *Shigella* germs spreading, wait at least two weeks after diarrhea ends to have sex.”



Written by [R. Cort Kirkwood](#) on March 3, 2023

We may assume from the emphasis that CDC has placed on the subject that homosexuals are behind a vast majority of XDR shigella infections.

For those wondering what “other men who have sex with men” means, CDC explains that the phrase is “used in CDC surveillance systems because it indicates men who engage in behaviors that may transmit Shigella infection, rather than how someone identifies their sexuality.”

Right.

Homosexuals and Monkeypox

The shigellosis outbreak is the second serious disease outbreak within a year caused by rampant homosexual activity. The monkeypox contagion began last year.

As [CDC reported](#), “at this time, data suggest that gay, bisexual, and other men who have sex with men make up the majority of cases in the current mpox outbreak.”

The [World Health Organization renamed](#) the disease mpox because of “racist and stigmatizing language online.”

Such is the association of the disease with homosexual sodomy that President Joe Biden picked a flaming leather fetishist who looks like a member of the Village People, [Demetre Daskalakis](#), as monkeypox czar.

Meet Dr. Demetre Daskalakis, Biden’s pick for White House Monkeypox response deputy coordinator.

Daskalakis is a gay pentagram wearing radical leftist- looks like he has all the qualifications necessary to be part of the Biden administration ☐☐ pic.twitter.com/c7gAudJZSY

— The Conservative Read☐☐ (@theconread) [September 8, 2022](#)

H/T: [Ace of Spades](#)



Subscribe to the New American

Get exclusive digital access to the most informative, non-partisan truthful news source for patriotic Americans!

Discover a refreshing blend of time-honored values, principles and insightful perspectives within the pages of "The New American" magazine. Delve into a world where tradition is the foundation, and exploration knows no bounds.

From politics and finance to foreign affairs, environment, culture, and technology, we bring you an unparalleled array of topics that matter most.



What's Included?

- 24 Issues Per Year
- Optional Print Edition
- Digital Edition Access
- Exclusive Subscriber Content
- Audio provided for all articles
- Unlimited access to past issues
- Coming Soon! Ad FREE
- 60-Day money back guarantee!
- Cancel anytime.

Subscribe